

Physician-led - Patient-centered Infusion-core

Please fax completed form, insurance card, and clinical documentation to: FAX: 410-220-2553

RITUXIMAB ORDER GPA/MPA	☐ New Start ☐ M	aintenance: Last Dose Given
Referring Office:	Contact Name:	Date:
Direct Phone for Contact:		Fax:
Patient Name:		DOB:
Allergies □ NKDA □ Allergies:		
	F	Height:
Weight:		
Indication: ☐ M31.30 Granulomatosis w/ Polyangiitis ☐ M31.7 Microscopic Polyangiitis (MPA) ☐ Other		
DRUG: Rituxan Truxima Riabni Ruxience ☐ Rituximab-per insurance preferred ☐ Rituxan ☐ Truxima (rituximab-abbs) ☐ Riabni (rituximab-arrx) ☐ Ruxience (rituximab-pvvr)		
INDUCTION DOSES: ☐ 375mg/m² every week X 4 weeks ☐ 1000mg IV at day 0 and 15 (approximately) ☐ Other	□ 500m	IANCE DOSES: g IV at day 0 and 15 nately) 500mg IV every

PREMEDICATION ORDERS: antihistamine, acetaminophen and 100mg methy the PI ☐ Acetaminophen po: ☐ 1000mg ☐ 500mg 30 min prior to infus ☐ Diphenhydramine: ☐ 25mg PO ☐ 50mg PO ☐ 25mg IVP 30 min pr☐ Solu-Medrol: ☐ 62.5mg IVP ☐ 100mg IVP ☐ Other 30 min pr☐ Other	rion. rior to infusion. prior to infusion.
Prescriber Name: Title:	
NPI: DEA:	
Prescriber Signature: Date of	Order:
L L	
Referrals will not be processed until we receive <u>ALL</u> the following:	
☐ Face Sheet / Patient Demographics	
☐ Insurance card(s) — copy of front & back	t & failed therapy
\square Last 2 clinic notes pertaining to referring diagnosis (include ALL pasoutcomes) Most Recent Labs (within last 4-8 weeks) – Required:	ε α raneu therapy
☐ CBC ☐ CMP ☐ TB ☐ Hep B Other:	